

**MATHEMATICS**

- 1.** Pairs of dice is thrown 5 times, if sum 5 is considered success than find probability of getting 4 success:

Ans. $\frac{40}{95}$

Sol. favour outcomes $\{(1,4), (4,1), (2,3), (3,2)\}$

$$\text{Success probability} = \frac{4}{36} = \frac{1}{9}$$

$$\text{Probability of getting 4 success} = {}^5C_4 \left(\frac{1}{9}\right)^4 \frac{8}{9} = \frac{5 \times 8}{95} = \frac{40}{95}$$

- 2.** Find coefficient of x^{18} in $\left(x^4 - \frac{1}{x^3}\right)^{15}$.

Ans. ${}^{15}C_6$

Sol. $T_{r+1} = {}^{15}C_r x^{60-4r} \times (-1)^r \times x^{-3r}$

$$60 - 7r = 18$$

$$r = 6$$

$$\text{Coefficient} = {}^{15}C_6$$

- 3.** Sum of first 20 terms of the sequence 5, 11, 19, 29,

Ans. 3520

Sol. $T_n = an^2 + bn + c = 5$

$$T_1 = a + b + c = 5 \quad \dots\dots(1)$$

$$T_2 = 4a + 2b + c = 11 \quad \dots\dots(2)$$

$$T_3 = 9a + 3b + c = 19 \quad \dots\dots(3)$$

$$(2) - (1) \Rightarrow 3a + b = 6$$

$$(3) - (2) \Rightarrow 5a + b = 8$$

$$a = 1, b = 3, c = 1$$

$$T_n = n^2 + 3n + 1$$

$$S_n = \frac{n(n+1)(2n+1)}{6} + \frac{3n(n+1)}{2} + n$$

$$S_{20} = \frac{20 \times 21 \times 41}{6} + \frac{3}{2} \times 20 \times 21 + 20$$

$$= 70 \times 41 + 630 + 20 = 3520$$

- 4.** $(p \rightarrow q) \wedge (r \rightarrow q)$ is equivalent to

$$(1) \sim p \wedge \sim r \quad (2) q \vee (\sim p \wedge \sim r) \quad (3) q \vee (\sim p \vee \sim r) \quad (4) \text{None of these}$$

Ans. (3)

Sol. $(\sim p \vee q) \wedge (\sim R \vee Q)$

$$= q \vee (\sim p \wedge \sim R)$$

Ans. (B)

-
5. If $2x^y + 3y^x = 20$ then find $\frac{dy}{dx}$ at $(2, 2)$

Ans. $\frac{-3}{5} \ln 2$

Sol. $2x^y + 3y^x = 20$

$$\Rightarrow 2 \cdot y \cdot x^{y-1} \cdot \frac{dy}{dx} + 3 \cdot y^x \left(\ln y + \frac{x}{y} \cdot \frac{dy}{dx} \right) = 0$$

$$\Rightarrow 2 \cdot \frac{dy}{dx} \cdot y \cdot x^{y-1} = (-3y^x \cdot \ln y)$$

$$\Rightarrow \frac{dy}{dx} [2y \cdot x^{y-1} + 3xy^{x-1}] = -3y^x \cdot \ln y$$

at $x = 2, y = 2$

$$\Rightarrow \frac{dy}{dx} [2 \times 2 \times 2^1 + 3 \times 2 \times 2^1] = -3 \times 4 \times \ln 2$$

$$\Rightarrow \frac{dy}{dx} [5] = -3 \ln 2 \Rightarrow \frac{dy}{dx} = \frac{-3}{5} \ln 2$$

6. For two groups of 15 sizes each, mean and variance of first group is 12, 14 respectively, and second group has mean 14 and variance σ^2 . If combined variance is 13 then find variance of second group?

(1) 9

(2) 11

(3) 10

(4) 12

Ans. (3)

Sol. $\frac{\sum x_i}{15} - (12)^2 = 14$

$$\sum x_i^2 = 158 \times 15$$

$$\frac{\sum x_i}{15} = 12$$

$$\sum x_i = 180$$

$$\frac{\sum y_i}{15} = 14$$

$$\frac{\sum y_i}{15} = 14$$

$$\sum y_i = 15 \times 14$$

$$\text{Combined mean} = \frac{\sum x_i + \sum y_i}{30} = \frac{15 \times 12 + 15 \times 14}{30} = 13$$

$$\frac{\sum y_i^2 + \sum x_i^2}{30} - (13)^2 = 13$$

$$\frac{\sum y_i^2 + 158 \times 15}{30} = 13 \times 14$$

$$\sum y_i^2 = 30 \times 13 \times 14 - 158 \times 15$$

$$\sigma^2 = \frac{30 \times 13 \times 14 - 158 \times 15}{15} - (14)^2$$

$$\sigma^2 = 10$$

7. A matrix of 2×2 satisfies $A^2 - I = 0$. If trace (A) = a and $|A| = b$ then find $3a^2 + 4b^2$.

Ans. 4

Sol. Characteristic equation of 2×2 matrix

$$A^2 - \text{tr}(A) \cdot A + |A| \cdot I = 0$$

$$\text{tr}(A) = 0$$

$$|A| = 1$$

$$3a^2 + 4b^2 = 4 \times 1 = 4$$

8. Given $f(x) = [a + 13 \sin x]$. Find the points of non-differentiability of $f(x)$ in $(0, \pi)$. Given 'a' is an integer

Ans. 25

Sol. $f(x) = a + [13 \sin x]$

When $13 \sin x = \text{An integer} = I$ (Let)

But $0 < \sin x \leq 1$ hence, $13 \sin x \in (0, 13]$

In $x \in (0, \pi)$ number of points where $13 \sin x \in (0, 13] = 12 \times 2 + 1 = 25$

9. Find the number of ways of distributing 20 oranges among 3 children such that each gets atleast one orange

Ans. ${}^{19}C_2$

Sol. $x_1 + x_2 + x_3 = 20$

$$x_i \geq 1$$

$$x_1' + x_2' + x_3' = 17 \quad (\text{let } x_i' = x_i + 1)$$


$$x_i' \geq 0$$

$$17 + 3 - 1 = {}^{19}C_2$$

10. $|x^2 - 5|x| + 7| = P \quad (P \in \mathbb{I})$ find number of possible values of "P" such that equation have 4 solutions.

Ans. 6


Sol.



- 11.** AB is building of height 30 m with angle of depression from top of building A to P. and Q is 15° and 60° respectively. A point C is on the same level on building with point P, then find the area of rectangle PCBQ?

Ans. $600(\sqrt{3} - 1)$

Sol.



$$\frac{30}{BQ} = \sqrt{3} \Rightarrow BQ = \frac{30}{\sqrt{3}} = 10\sqrt{3}$$

$$\frac{AC}{10\sqrt{3}} = 2 - \sqrt{3}$$

$$AC = (2 - \sqrt{3})10\sqrt{3}$$

$$\text{Area} = 10\sqrt{3} [60 - 20\sqrt{3}] = 600(\sqrt{3} - 1)$$

- 12.** Given that $f(x) + 4f\left(\frac{1}{x}\right) = \frac{1}{x} + 3$, find the value of $\int_1^2 18f(x) dx$.

Ans. $10\ln 2 - 6$

$$5f(4) + 4f\left(\frac{1}{4}\right) = \left(\frac{1}{4} + 3\right) \times 5$$

$$5f\left(\frac{1}{x}\right) + 4f(x) = x + 3 \times 3$$

$$9f(x) = \frac{5}{x} - 4x + 3$$

$$f(x) = \frac{5}{9x} - \frac{4x}{9} + \frac{1}{3}$$

$$\int_1^2 18f(x) dx = \int_1^2 \left(\frac{1}{x} - 8x + 6 \right) dx$$

$$= \left[10\ln x - \frac{8x^2}{2} + 6x \right]_1^2$$

$$= 10\ln 2 - 6$$

- 13.** If the image of point P(1,2,3) about the plane $2x-y+3z=2$ is Q, then the area of triangle PQR.
Where coordinates of R is (4,10,12).

Ans. $\frac{\sqrt{1531}}{2}$

Sol. Image formula $\rightarrow \frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{2} = \frac{-2(ax_1 + by_1 + cz_1 d)}{a^2 + b^2 + c^2}$

$$\frac{x-1}{2} = \frac{x-y_1}{-1} = \frac{2-3}{3} = \frac{-2(2-2+9-2)}{4+1+9}$$

$$\Rightarrow x = -1$$

$$y = 3$$

$$z = 0$$

so, P(1, 2, 3)

Q(1, 2, 3)

$$\vec{PQ} = -2\mathbf{i} + \mathbf{j} - 3\mathbf{k}$$

R(4, 10, 12)


$$\vec{PR} = 3\mathbf{i} + 8\mathbf{j} - 3\mathbf{k}$$

$$\text{Area if } \Delta OQR = \frac{1}{2} |\vec{PQ} \times \vec{QR}|$$

$$= \frac{1}{2} [i(9+24) - j(-18+9) + k(-16-3)]$$

$$= \frac{1}{2} [3\hat{i} + 9\hat{j} - 19\hat{k}]$$

$$= \frac{\sqrt{33^2 + 9^2 + 19^2}}{2} = \frac{\sqrt{1531}}{2}$$



- 14.** The sum of roots of $|x^2 - 8x + 15| - 2x + 7 = 0$

(1) $11 + \sqrt{3}$

(2) $11 - \sqrt{3}$

(3) $9 + \sqrt{3}$

(4) $9 - \sqrt{3}$

Ans. (3)

Sol. (1) $\rightarrow x \in (-\infty, 3] \cup [5, \infty)$

$$(x-3)(x-5) - 2x + 7 = 0$$

$$\Rightarrow x^2 - 8x + 15 - 2x + 7 = 0$$

$$\Rightarrow x^2 - 10x + 22 = 0 \rightarrow x = \frac{10 \pm \sqrt{100 - 88}}{2}$$

$$x = 5 \pm \sqrt{3}$$

$$x = 5 + \sqrt{3}$$

(2) $\rightarrow x \in (3, 5)$

$$-(x-3)(x-5) - 2x + 7 = 0$$

$$-x^2 + 8x + 15 - 2x + 7 = 0$$

$$\Rightarrow (x^2 - 6x + 8) = 0$$

$$x = 4$$

$$\text{Sum of roots} = 5 + \sqrt{3} + 4$$

$$= 9 + \sqrt{3}$$

- 15.** let $\vec{a} = 2\hat{i} + 3\hat{j} + 4\hat{k}$, $\vec{b} = \hat{i} - 2\hat{j} - 2\hat{k}$, $\vec{c} = -\hat{i} + 4\hat{j} + 3\hat{k}$ and \vec{d} is vector perpendicular to both \vec{b} and \vec{c} and $\vec{a} \cdot \vec{d} = 18$, then $|\vec{a} \times \vec{d}|^2$ is

(1) 640 (2) 720 (3) 680 (4) 760

Ans. (2)

Sol. $d = \lambda(\bar{b} \times \bar{c})$

$$\lambda \begin{vmatrix} i & j & k \\ 1 & -2 & -2 \\ -1 & 4 & 3 \end{vmatrix} = \lambda(\hat{i}(2) - \hat{j}(1) + \hat{k}(2))$$

$$\Rightarrow \lambda(2\hat{i} - \hat{j} + 2\hat{k})$$

$$(\bar{a} \times \bar{d}) = (2 \ 3 \ 4) \times (4 \ -2 \ 4)$$

$$\begin{vmatrix} i & j & k \\ 2 & 3 & 4 \\ 4 & -2 & 4 \end{vmatrix} = \hat{i}(20) - \hat{j}(-8) + \hat{k}(-16)$$

$$20\hat{i} + 8\hat{j} - 16\hat{k}$$

$$400 + 64 + 256$$

$$= 720$$

- 16.** Let $a_1, a_2, a_3 \dots, a_n$ are in arithmetic progression having common difference d. The value of

$$\lim_{n \rightarrow \infty} \sqrt{\frac{d}{n}} \left(\frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_3}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} \right) \text{ is:}$$

Ans. 1

Sol. $\lim_{n \rightarrow \infty} \sqrt{\frac{d}{n}} \left(\frac{1}{\sqrt{a_1} + \sqrt{a_2}} + \frac{1}{\sqrt{a_2} + \sqrt{a_5}} + \dots + \frac{1}{\sqrt{a_{n-1}} + \sqrt{a_n}} \right)$ is

$$\frac{(\sqrt{a_1} - \sqrt{a_2})}{(\sqrt{a_1} - \sqrt{a_2})(\sqrt{a_1} + \sqrt{a_2})} = \begin{pmatrix} \sqrt{a_1} - \sqrt{a_2} \\ -d \end{pmatrix}$$

$$\lim_{n \rightarrow \infty} \sqrt{\frac{d}{n}} \cdot \frac{1}{d} \left(\sqrt{a_2} - \sqrt{a_1} - \sqrt{a_2} - \sqrt{a_1} - \dots - \sqrt{a_n} - \sqrt{a_{n-1}} \right)$$

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{nd}} \frac{(\sqrt{a_n} - \sqrt{a_1})(\sqrt{a_n} + \sqrt{a_1})}{\sqrt{a_n} + \sqrt{a_1}}$$

$$\lim_{n \rightarrow \infty} \frac{1}{\sqrt{n}\sqrt{d}} \frac{a_1 + (n-1)d - a_1}{\sqrt{a_n} - \sqrt{a_1}}$$

$$\lim_{n \rightarrow \infty} \frac{(n-1)\sqrt{d}}{\sqrt{n}\sqrt{a_1 + (n-1)d} + \sqrt{a_1}}$$

$$\lim_{n \rightarrow \infty} \frac{\left(1 - \frac{1}{n}\right)\sqrt{d}}{\sqrt{\frac{a_1(n-1)d}{\sqrt{n}} + \frac{\sqrt{a_1}}{\sqrt{n}}}} = \frac{\sqrt{d}}{\sqrt{d} + 0} = 1$$

$$\sqrt{\frac{a_1}{n} + \frac{(n-1)d}{n}}$$



- 17.** The ratio of terms of 5th term from beginning and 5th term from end is $\sqrt{6} : 1$ in $\left(\frac{\frac{1}{2^4}}{\frac{1}{3^4}} + 1 \right)^n$ the value of n is:

Ans. 10

Sol. $T_5 = {}^nC_4 \left(\frac{1}{2^4} \right)^{n-4} \left(\frac{1}{\frac{1}{3^4}} \right)^4$

5th term from last = ${}^nC_4 \left(\frac{1}{\frac{1}{3^4}} \right)^{n-4} \left(\frac{1}{2^4} \right)^4$

$$\frac{2^{\frac{n-4}{4}}}{3} \cdot \frac{3^{\frac{n-4}{4}}}{2} = \frac{\sqrt{6}}{1}$$

$$3^{\frac{n-8}{4}} \cdot 2^{\frac{n-8}{4}} = 6^{\frac{1}{2}}$$

$$\frac{n-8}{4} = \frac{1}{2} \Rightarrow 2n - 16 = 4$$

$$2n = 20$$

$$n = 10$$

- 18.** If ${}^{2n}C_3 : {}^nC_3 = 10$, then $\frac{n^2 + 3n}{n^2 - 3n + 4}$ is equal to

Ans. 2

Sol. if ${}^{2n}C_3 : {}^nC_3 = 10$, then $\frac{n^2 + 3n}{n^2 - 3n + 4}$ is equal to.....

$$\Rightarrow \frac{(2n)(2n-1)(2n-2)}{n(n-1)(n-2)} = 10$$

$$2.2 \frac{(2n-1)}{(n-2)} = 10 \Rightarrow 4n-2 = 5n-10$$

$$\boxed{n = 8}$$

$$\frac{64+24}{64-24+4} = \frac{88}{44} = 2$$

- 19.** The integration $\int \frac{x^2(x \sec^2 x + \tan x)}{(x \tan x + 1)^2} dx$ is

(1) $\frac{x}{x \tan x + 1} + \log|x \sin x + \cos x| + C$

(2) $\frac{x}{x \tan x + 1} - \log|x \sin x + \cos x| + C$

(3) $-\frac{x^2}{x \tan x + 1} + 2 \log|x \sin x + \cos x| + C$

(4) $\frac{x^2}{x \tan x + 1} + 2 \log|x \sin x + \cos x| + C$

Ans. (3)

Sol. $\int \frac{x^2(x \sec^2 x + \tan x)}{(x \tan x + 1)^2} dx$

Apply by parts and substituting $(x \tan x + 1) = t$ to solve the integral we get



$$\text{We get } I = \frac{-x^2}{x \tan x + 1} + \int \frac{2x \cos(x)}{x \sin x + \cos x} dx$$

$$\text{Now substituting } x \sin x + \cos x = t \text{ we get } I = \frac{-x^2}{x \tan(x) + 1} + 2 \ln(|x \sin x + \cos x|) + C$$